

## ICE Checklist #2: Parenteral Medication Preparation and Administration

Certification Number: \_\_\_\_\_

Observation 1: Shift # \_\_\_\_\_ Staff Type \_\_\_\_\_ Isolation Y ☐ N ☐ Visible From Nursing Station Y ☐ N ☐

Observation 2: Shift # \_\_\_\_\_ Staff Type \_\_\_\_\_ Isolation Y ☐ N ☐ Visible From Nursing Station Y ☐ N ☐

### Hand hygiene before preparing medications

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Medications prepared in a clean area, on a clean surface, away from dialysis stations

**Exception:** Drawing saline syringes at dialysis station from patient's own clean saline bag, using aseptic technique

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Assemble supplies: sterile syringes, 70% alcohol swabs or other antiseptic, medication vials

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Single-dose vials used for one patient only and discarded (punctured only one time)

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Multiple-dose vials only entered with a new, empty sterile syringe and needle and discarded within 28 days unless manufacturer specifies a different (shorter or longer) date for that opened vial (see Information Sheet #4)

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Open one vial of each medication at a time; wipe stopper with alcohol or other antiseptic; withdraw medication into sterile syringe. May prepare meds for multiple patients at one time, but must administer to one patient at a time, leaving remainder of drawn meds in clean preparation area

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Label syringes that are predrawn and not immediately administered with patient name, medication, dose, time drawn; take only individual patient's medications to dialysis station

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Hand hygiene

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Don clean gloves; wipe injection port (or patient's skin if subcutaneous or intramuscular injection) with antiseptic (e.g., chlorhexidine, povidone iodine, iodophor, or 70% alcohol); inject medication

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Discard syringe into Sharps container at point of use; remove gloves

**Exception:** If using a needleless system with no attached needle, disposal in Sharps not necessary

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Hand hygiene

Obs 1: Met ☐ Not Met ☐

Obs 2: Met ☐ Not Met ☐

### Observation 1 notes:

### Observation 2 notes: